

WO 01/17344

PCT/CZ00/00064

1/12

A photography of the seed producing ducts after repeated (5-times 8 Gy dose) irradiation of testicles of an acceptor cock. In the sperm producing ducts only the Sertoli's cells are visible, the germline spermatogonial cells are not visible (dyed by Pas, enlarged 400-times).





WO 01/17344


2/12

PCT/CZ00/00064

Minorca black
(ii,EE,b/b)




suspension of
spermatogonial
cells




Leghorn white
(II)

X



Leghorn buff
(ii,ee,B/-)



(ii,Ee,B/b)



(ii,Ee,b/-)


Fig. 2

WO 01/17344

3/12

PCT/CZ00/00064

Fig. 3




WO 01/17344

4/12

PCT/CZ00/00064

Fig. 4




WO 01/17344

5/12

PCT/CZ00/00064

Fig. 5




WO 01/17344

6/12

PCT/CZ00/00064

Fig. 6




WO 01/17344

7/12

PCT/CZ00/00064

Fig. 7




WO 01/17344

8/12

PCT/CZ00/00064

Fig. 8




WO 01/17344

9/12

PCT/CZ00/00064

Fig. 9




WO 01/17344

10/12

PCT/CZ00/00064

Fig. 10




WO 01/17344

11/12

PCT/CZ00/00064

Fig. 11



WO 01/17344

PCT/CZ00/00064

12/12

Fig. 12

